

# Base map compiled by Illinois State Geological Survey from digital data (2009 TIGER/Line Shapefiles) provided by the United States Census Bureau.

North American Datum of 1927 (NAD 27) Projection: Transverse Mercator 10,000-foot ticks: Illinois State Plane Coordinate system, west zone (Transverse Mercator) 1,000-meter ticks: Universal Transverse Mercator grid system, zone 16

#### **Recommended citation:**

Kolata, D.R., 2012, Bedrock Geology of Grand Detour Quadrangle, Ogle and Lee Counties, Illinois: Illinois State Geological Survey, Illinois Geologic Quadrangle Map, IGQ Grand Detour-BG, 2 sheets, 1:24,000.



© 2012 University of Illinois Board of Trustees. All rights reserved. For permission information, contact the Illinois State Geological Survey.

#### Geology based on field work by D.R. Kolata, 2010–2011.

Digital cartography by Jane E. Johnshoy Domier, Trisha S. Rentschler, and Coy E. Potts, Illinois State Geological Survey.

This research was supported in part by the U.S. Geological Survey National Cooperative Geologic Mapping Program (STATEMAP) under USGS award number G10AC00418. The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Government.

The Illinois State Geological Survey and the University of Illinois make no guarantee, expressed or implied, regarding the correctness of the interpretations presented in this document and accept no liability for the consequences of decisions made by others on the basis of the information presented here. The geologic interpretations are based on data that may vary with respect to accuracy of geographic location, the type and quantity of data available at each location, and the scientific and technical qualifications of the data sources. Maps or cross sections in this document are not meant to be enlarged.





Prairie Research Institute Illinois State Geological Survey 615 East Peabody Drive Champaign, Illinois 61820-6964 (217) 244-2414 http://www.isgs.illinois.edu





Light-duty road, hard or Secondary highway, hard surface improved surface \_\_\_\_\_

State Route

# IGQ Grand Detour-BG Sheet 1 of 2

|   | SYSTEM          | SERIES                              | GROUP       | FORMATION<br>or<br>MEMBER                                       | GRAPHIC COLUMN | THICKNESS<br>(feet) |         | UNIT |
|---|-----------------|-------------------------------------|-------------|---|----------------|---------------------|---------|------|
|   | QUATER-<br>NARY | HOLOCENE<br>AND<br>PLEISTO-<br>CENE |             | Surficial<br>deposits   |                | C                   | -200    | А    |
|   |                 |                                     | Galena      | Wise Lake   |                | 0-220               | 0–80    | В    |
|   |                 |                                     |             | Haldane K-bentonite -<br>Dunleith                               |                |                     | 120–130 | С    |
|   |                 |                                     |             | Guttenberg  |                |                     | 0–6     | D    |
|   | ORDOVICIAN      | MOHAWKIAN                           | Platteville | Quimbys Mill  |                |                     | 10–15   | Е    |
|   |                 |                                     |             | Nachusa   |                |                     | 15–25   | F    |
|   |                 |                                     |             | Grand Detour  |                | 110–130             | 30–40   | G    |
|   |                 |                                     |             | Mifflin   |                |                     | 15–25   | н    |
| Goologia Fosturas                                     |                 |                                     |             | Pecatonica  |                |                     | 20–35   | I    |
| K-bentonite bed<br><i>Fisherites (Receptacutites)</i> |                 |                                     | Ancell      | Harmony Hill Sh<br>Loughridge Ss<br>Daysville Dol<br>Kingdom Ss |                | 175–435             | 25–35   | J    |
| ت Hardground surface                                  |                 |                                     |             | St Potor  |                |                     | 100     |      |

A Quaternary Surficial Deposits Clay, silt, sand, gravel, and boulders deposited primarily as alluvium along river and stream valleys, rock debris below cliffs, glacial till and windblown silt on the uplands, and local lacustrine deposits. These units are not shown on map.

**B** Wise Lake Formation Dolomite, light gray in fresh exposures, weathers to light buff or yellowish gray; pure, vesicular, vuggy in weathered outcrops, medium grained; compared with the underlying Dunleith Formation, the unit is non-cherty except in the lower 2 to 3 feet; weathers to 6- to 12-inch-thick even beds. Relict bioclastic textures indicate that the initial carbonate deposit consisted of lime mudstone and wackestone with a few interspersed beds of grainstone. A 1- to 2-inch-thick volcanic ash bed (Dygerts K-bentonite) is present locally near the base of the formation (Willman and Kolata 1978). Fossils consist mainly of dolomitized molds and casts of whole and broken specimens. Some of the most common fossils include the brachiopod Rafinesquina; gastropods Hormotoma and Liospira; coral Streptelasma; trilobite Illaenus; and blue-green algae Fisherites (Receptaculites). The Wise Lake occurs at the bedrock surface in the western parts of the quadrangle and is well exposed along the bluffs of Pine Creek in and near White Pines State Park. The Wise Lake is gradational with the underlying Dunleith Formation.

**C Dunleith Formation** Dolomite, light gray in fresh exposures, weathers to light buff or yellowish gray; argillaceous and slightly shaly at the base grading upward to more pure dolomite. Vuggy in weathered outcrops, medium grained, vesicular; weathers to mainly 2- to 8-inch-thick beds that are even to wavy and nodular. Contains white chert nodules, lenses, and thin beds. The amount of chert is variable but generally less than 5% except for a few beds 5 to 10 feet thick, which in places are up to 15 or 20% of the rock volume. Hardground omission surfaces are present throughout but are particularly abundant in the lower part of the formation. Some surfaces are widespread and potentially useful for detailed correlation of strata. A 1- to 2-inchthick volcanic ash bed (Haldane K-bentonite) is present locally near the top of the formation (Willman and Kolata 1978). Fossils are locally abundant and consist of molds and casts within the beds and dolomitized shelly fossils on some bedding planes. Trepostome bryozoans are particularly abundant in the lower part of the formation. Other common fossils include the brachiopods *Rafinesquina*, *Dalmanella*, and Sowerbyella; blue-green algae Fisherites (Receptaculites) is common in an 8-foot zone beginning about 20 feet above the base. The Dunleith is present at the bedrock surface in the western parts of the quadrangle with notable outcrops along the western bluffs of the Rock River in Sec. 17, 18, 20, and 21, T22N, R9E. The base is commonly marked by a hardground surface.

**D** Guttenberg Formation Dolomite, light pinkish tan in fresh exposures; argillaceous, vesicular, medium grained; 2- to 4-inch-thick wavy, nodular beds separated by thin beds of dark reddish brown carbonaceous shale as much as an inch thick but mostly less than 1/4 inch. White to medium gray chert nodules are present locally. Fossils are abundant and diverse; commonly molds and casts of gastropods, cephalopods, and bivalves are present within beds; bedding surfaces are commonly covered with well-preserved shelly fossils including various echinoderms, bryozoans, and brachiopods Rafinesquina, Dalmanella and Sowerbyella. A K-bentonite bed up to 2 inches thick is locally present at the base of the formation. It is present in the rock core obtained near the intersection of Lowell Park Road and Penn Corners Road (SE NE NE Sec. 31, T23N, R9E; IP number 121412600200; ISGS core C15474). The Guttenberg is thin (6 feet or less) to absent in the Grand Detour Quadrangle. Outcrops are restricted to the western bluffs of the Rock River in Sec. 17, 18, 20, and 21, T22N, R9E and southwest parts of Sec.16, T23N, R9E. The top and bottom of the ormation are marked by prominent naroground surfaces.

**E** Quimby's Mill Formation Dolomite, very light gray in fresh exposures; argillaceous to relatively pure, dense, fine to very fine grained; 2- to 8-inch-thick, even, well-defined beds with smooth wavy bedding planes separated by light greenish gray shale partings mainly less than 1/8 inch thick; locally contains white chert nodules. Relict bioclastic textures in the chert indicate that the initial carbonate deposit was lime mudstone. Lenticular cavities 1/2 to 1 inch across, up to 1/8 inch wide, and vertically oriented characterize the upper part of the formation. The formation has been mined for flagstone in the southwest part of Sec. 16, T23N, R9E. The fauna is sparse and low in diversity. The top of the formation is marked by a prominent hardground omission surface that is pocked with irregular solution cavities up to 2 inches across and 1 to 6 inches deep that are encrusted with iron-rich minerals. The base is gradational with the underlying Nachusa Formation.

**F** Nachusa Formation Dolomite, light gray in fresh exposures, weathers to light buff or yellowish gray; pure to slightly argillaceous, vesicular, vuggy in weathered outcrops, fine- to medium-grained, 4- to 8-inch-thick beds; contains white to medium gray chert nodules. Relict bioclastic textures in the chert indicate that the initial carbonate deposit was lime mudstone. The formation is readily divisible into three members in the Grand Detour Quadrangle based on dolomite purity. The relatively pure Eldena (oldest) and Everett Members are separated by the argillaceous dolomite of the Elm Member (not shown). The shelly fossils are sparse and low in diversity; however, there is a conspicuous abundance of trace fossils assignable to the genus Chon*drites* that occur on bedding planes throughout the formation. The base of the formation is commonly marked by an abrupt change from relatively pure dolomite to the underlying clay- and silt-rich dolomite of the Grand Detour Formation.

**G** Grand Detour Formation Dolomite and limestone, light gray in fresh exposures, weathers to light buff or yellowish gray; pure to argillaceous, fine grained; contains white to medium gray chert nodules. Upper part of the formation is argillaceous and consists of wavy, nodular beds mostly between 3 and 6 inches thick that are separated by medium brownish gray shale partings up to 1 inch thick. These beds are characterized by medium to dark gray spots approximately <sup>1</sup>/<sub>8</sub> inch in diameter. The lower part of formation is relatively pure with even beds that are mostly 4 to 10 inches thick and contain white to medium gray chert nodules. Limestone is present in the southern parts of the quadrangle, most notably in the cement quarries in Sec. 16, T22N, R9E. One or more hardground omission surfaces are present in the lower part of the formation. A 2- to 4-inch-thick bed of upward fining grainstone is also present near the base. The Grand Detour contains diverse and locally abundant fossil fauna including brachiopods Opikina, Campylorthis, Hesperorthis, Rostricellula, and Strophomena; gastropods Lophospira, Clathrospira, Ectomaria, Phragmolites, Tetranota, and Maclurites; cephalopods Richardsondoceras and Endoceras; trilobites Thaleops, Ceraurus, Encrinurus, and Isotelus; corals *Foerstephyllum* and *Streptelasma*; and crinoids *Cupulocrinus* and Ablutoglyptocrinus. The lower 2 feet of the formation contains a locally abundant lithistid fossil sponge fauna including Anthaspidella and *Zittelella*. The base is marked by a hardground surface.

**H** Mifflin Formation Dolomite and limestone, light gray in fresh exposures, weathers to light buff or yellowish gray; argillaceous, fine to very fine-grained lime mudstone and wackestone; 1- to 4-inch-thick wavy, nodular beds separated by beds of greenish gray shale as much as an inch thick, but mostly less than 1/2 inch. A 2- to 4-inch-thick bed of fining upward grainstone is present near the top of the formation. The abundance of shaly partings and nodular bedding differentiate the Mifflin from the thicker-bedded, relatively pure formations above and below. Calcium-rich, low magnesium limestone is present in the southern half of the quadrangle where it is mined for the manufacture of cement. The formation is entirely dolomite in exposures along the Burlington-Northern Railroad tracks in Sec. 7, T23N, R10E. A thin K-

bentonite bed (not shown) has been observed locally at the top of the formation (Willman and Kolata 1978). The Mifflin is the most fossiliferous unit in the Platteville Group. Bedding planes, particularly in the limestone facies, are typically covered with well-preserved disarticulated or whole invertebrate fossils including brachiopods, bryozoans, ostracodes, corals, trilobites, and echinoderms. An abundant and diverse fossil molluscan fauna is commonly present within beds. Some of the more common fossils include the brachiopods *Opikina*, *Campylorthis*, Doleroides, Hesperorthis, Protozyga, Rostricellula, and Strophomena; gastropods Lophospira, Clathrospira, Ectomaria, Phragmolites, Tetranota, Subulites, and Maclurites; cephalopods Richardsondoceras, Whitfieldoceras, and Endoceras; clam Vanuxemia; trilobites Thaleops, Ceraurus, Encrinurus, and Isotelus; corals Foerstephyllum and Streptelasma; and crinoids Cupulocrinus, Cremacrinus, and Ablutoglypto*crinus*. The base of the formation is marked by a prominent ferruginous hardground surface.

**I Pecatonica Formation** Dolomite and limestone, light gray in fresh exposures, weathers to light buff or yellowish gray; pure to slightly argillaceous, fine grained, dense; 2- to 8-inch-thick even, well-defined beds with light greenish gray shale partings mainly less than 1/8 inch thick; a few beds near the middle of the unit are wavy and nodular; contains white to medium gray chert nodules. The lower half of the formation contains St. Peter Sandstone-like quartz sand grains that are well-rounded and frosted and increase in abundance downward. Phosphatic grains and nodules are locally present in the lower 5 feet. The Pecatonica contains dolomite-mottled limestone near the town of Grand Detour and adjacent areas south and east of the Rock River. A deeply sculpted hardground omission surface is present at the top of the unit, and two or three are present in the lower 8 feet. Limestone is exposed in the cement quarries north of Dixon, Illinois (Sec. 22, T22N, R9E). The Pecatonica is less fossiliferous than the overlying Mifflin Formation, but a few beds in the lower part of the unit contain brachiopods, bryozoans, ostracodes, corals, trilobites, and echinoderms. The base is marked by an abrupt change from dolomite to the underlying shale.

J Glenwood Formation Shale, sandstone, siltstone, and dolomite. Four stratigraphic members can be recognized in the area of the Grand Detour Quadrangle. In ascending order these are the Kingdom Sandstone Member consisting of sandstone that is mainly greenish gray, silty, argillaceous, and pyritic; the Daysville Dolomite Member consisting mainly of dolomite that is argillaceous, silty, sandy, greenish gray, and chalky to dense; the Loughridge Sandstone Member composed mainly of sandstone that is silty, argillaceous, greenish gray or light brown, very fine to coarse grained, the mixed grain sizes producing a characteristic bimodal texture; and the Harmony Hill Shale Member consisting of shale that is medium to dark greenish gray or dark maroon, pyritic, partly silty and sandy, friable, and thinly laminated. The upper part of the Harmony Hill Shale contains phosphatic pellets and small nodules. The Glenwood Formation generally forms a sharp contact with the overlying dolomite of the Pecatonica Formation. The Glenwood is exposed in the upper reaches of the ravines in the western bluffs of the Rock River in Castle Rock State Park and in a quarry situated near the center of Sec. 18, T23N, R10E. Scolecodonts have been reported from the Glenwood (Templeton and Willman 1963), but the unit is largely unfossiliferous. The base is marked by an abrupt change to clean, white sandstone. Locally, the contact is an angular unconformity (fig. 2).

**K** St. Peter Sandstone Sandstone, pure guartz sand that is very light gray to white, very fine to coarse grained, well-rounded, frosted, and sorted, friable, and thick bedded. The St. Peter is well exposed in the ravines in the western bluffs of Rock River in Castle Rock State Park and is mined for industrial sand in a quarry situated near the center of Sec. 18, T23N, R10E. The base is covered. Local water wells indicate thicknesses ranging from 100 to 350 feet.





# INTRODUCTION

The Grand Detour Quadrangle is situated in the southwestern part of Ogle and northernmost part of Lee Counties, Illinois. The name is derived from the small town of Grand Detour, which sits at the eastern end of a major bend in the Rock River. The river flows north past the town, rather than its normal southwestern course; hence, the name Grand Detour.

The quadrangle lies in the Rock River Hill Country of the Central Lowlands Province. The upland areas are mostly cultivated for corn and soybeans. Limestone is mined for cement production and aggregate use in the southernmost parts of the quadrangle, and industrial sand is mined in the northeastern region. Other activities include public recreation in White Pines and Castle Rock State Parks as well as a moderate amount of residential and commercial developments.

Compilation of this map is based on an examination of bedrock exposures in quarries, road excavations, railroad cuts, and natural exposures along streams and waterways. Subsurface information was obtained from water well records, drill cuttings, and exploratory borings. In 2011 a continuous cored test hole was drilled to 260 feet by the Illinois State Geologic Survey (ISGS) in an abandoned quarry 5 miles north of Dixon, Illinois in Ogle County (IP 121412600200; ISGS core C15474 SE NE NE Sec. 31, T23N, R9E; latitude 41° 56' 30.45381" N and longitude 89° 29' 39.93540" W; height 775.3 feet). Wireline logs were also obtained from this key reference boring and filed in the Geological Records Unit of the ISGS. The uppermost beds exposed in the quarry are of the lower Wise Lake Formation, which overlie approximately 20 feet of the Dunleith Formation. The cored interval includes the remainder of the Dunleith Formation and the Quimbys Mill, Nachusa, Grand Detour, Mifflin, and Pecatonica Formations of the Platteville Group, the Glenwood Formation, and the uppermost St. Peter Sandstone. Field notes made by previous geologists and filed in the ISGS Library were another valuable source of information.

The bedrock in Lee and Ogle Counties was illustrated very generally on early statewide geologic maps (Worthen 1875, Weller 1906); however, the first concerted investigations of the Grand Detour area were published by Knappen (1926) in his report of the Dixon 15-minute Quadrangle, which includes the area covered by the Grand Detour 7.5-minute Quadrangle. A map showing the bedrock geology of Lee County, including the southern parts of the Grand Detour Quadrangle, was published by McGarry (1999). Stratigraphic and structural investigations in the Grand Detour area include those by Templeton and Willman (1952, 1963), Kolata and Buschbach (1976), Willman and Kolata (1978), and Kolata et al. (1978).

## **STRATIGRAPHY**

Throughout most of the uplands in the Grand Detour Quadrangle, the bedrock is covered by less than 25 feet of glacial drift, and, in many areas between Pine Creek and Rock River, bedrock occurs at the surface. Quaternary valley train deposits range from 0 to 50 feet thick (Piskin and Bergström 1975). The stratigraphic units that are known to occur at the bedrock surface in the quadrangle include the Upper Ordovician Ancell, Platteville, and Galena Groups. All were deposited in intertidal to subtidal marine environments. The Glenwood subdivisions that are present include the Kingdom Sandstone, Daysville Dolomite, Loughridge Sandstone, and Harmony Hill Shale Members. The Platteville and Galena Groups consist mainly of dolomite and local occurrences of limestone (Platteville only). These groups are further subdivided into members and formations based on relative amounts of disseminated clay and silt as well as the presence or absence of chert. Recognizable units within the Platteville include the Pecatonica, Mifflin, Grand Detour, Nachusa, and Quimbys Mill Formations. Galena rock units that are recognized in the Grand Detour Quadrangle include the Guttenberg, Dunleith, and Wise Lake Formations. Two or three K-bentonite beds are present in the Platteville and Galena rocks in this part of Illinois. These ash deposits originated from volcanic arcs situated along the southeastern margins of the continent in the vicinity of present-day North Carolina and South Carolina (Kolata et al. 1996). For all practical purposes, the ash desposits are isochronous.

Several hardground omission surfaces within the Galena and Platteville are widespread and persistent They are readily identified in outcrop and drill core and are useful in correlating this part of the strati-2 inches across. They penetrate the bedding planes to depths of 2 to 3 inches, are stained by iron-rich the more significant surfaces include (1) approximately 15 to 20 surfaces in the lower part of the Dun-

# STRUCTURAL GEOLOGY

150-

The Grand Detour Quadrangle lies along the northern margin of the Illinois Basin and is situated in a structurally complex area near the termination of the Sandwich Fault Zone, Plum River Fault Zone, and the projected trend of the La Salle Anticlinorium. Lying parallel to and south of the Sandwich Fault Zone in Lee and Ogle Counties is the Ashton Anticline (fig. 1). This broad anticline brings Middle Ordovician and Upper Cambrian rocks to the bedrock surface along the south side of the Sandwich Fault Zone just east of the Grand Detour Quadrangle and is responsible for the pronounced west-southwest dip within the quadrangle (see cross section A-A').



**Figure 1** The Sandwich Fault Zone and the Plum River Fault Zone.

## **Sandwich Fault Zone**

The Sandwich Fault zone extends in a northwesterly direction from near Manhattan, Will County, to central Ogle County, a distance of about 85 miles (Kolata et al. 1978). Outcrop and subsurface informa tion indicate that the fault zone is about 1/2 to 2 miles wide and is upthrown on the south along most of its extent. The fault zone has a maximum cumulative displacement of about 800 feet at its midpoint in southeastern De Kalb County.

One of the best exposures of the fault zone can be seen in the northeast corner of the Grand Detour Quadrangle at the Burlington Northern Railroad cut (NW NW SE and NE SW Sec. 7, T23N, R10E) and at outcrops in nearby quarries (SE SE Sec. 7, and NW SW Sec. 8, T23N, R10E) (Templeton and Willman 1952, Kolata et al. 1978). Here the Sandwich Fault Zone is characterized by about two dozen northwest-trending high-angle faults that bound small graben and horst blocks that displace strata ranging from the Pecatonica Formation to the Wise Lake Formation. Most of the faults have small displacements of 1 to 10, feet but one fault near the southwest end of the railroad cut may have as much as 130 feet of displacement. The faults project northwestward toward the Mt. Morris Quadrangle. Stratigraphic relations indicate that the faulting is post-Niagaran (middle Silurian) to pre-Illinoian (Pleistocene) in age and likely to be contemporaneous with the late Paleozoic deformation that formed many of the structural features in the midcontinent United States.

During recent mining operations near the center of Sec. 18 (T23N, R10E), a high wall (fig. 2) was cut through the Late Ordovician St. Peter Sandstone and overlying Glenwood Formation. This outcrop is remarkable because it reveals a previously unknown episode of Late Ordovician structural deformation in northern Illinois (Seid and Kolata 2011). The exposure is approximately 100 feet high and 900 feet long and extends in an east-west direction along a north-facing wall. The St. Peter Sandstone is folded into a broad north-northeast-trending syncline and anticline. Structurally tilted up to 20 degrees, the St Peter strata were subsequently erosionally truncated and covered by flat-lying shale, dolomite, and sandgraphic column. These planar surfaces are characterized by irregular solution cavities, mainly less than stone of the Glenwood Formation, forming a pronounced angular unconformity. Stratigraphic relations indicate that at least 100 feet of the St. Peter Sandstone was eroded from the crest of the anticline prior minerals, and are backfilled with sediment from the overlying unit (Kolata et al. 1998, 2001). Some of to deposition of the Glenwood. This site is situated along the northwestern termination of the Sandwich Fault Zone, which suggests that the folds are structurally associated with a zone of crustal weakness that leith Formation, (2) the top of the Guttenberg Formation, (3) the top of the Quimbys Mill Formation, (4) was episodically reactivated by tectonic forces during the Paleozoic. the top of the Mifflin Formation, (5) the top of the Pecatonica Formation, and (6) two or three surfaces in the lower 10 feet of the Pecatonica Formation.

#### **Small-Scale Domes, Folds and Faults**

Some of the most extensive bedrock outcrops in the Grand Detour Quadrangle are exposed in the openpit limestone mines north of Dixon, Illinois (S<sup>1</sup>/<sub>2</sub> Sec. 16, SW<sup>1</sup>/<sub>4</sub> 15, S<sup>1</sup>/<sub>2</sub> NE<sup>1</sup>/<sub>4</sub> Sec. 21, and S<sup>1</sup>/<sub>2</sub> Sec. 22, T22N, R9E). The common mining practice here has been to extract the Mifflin Formation for cement production leaving the hardground at the top of Pecatonica Formation exposed in wide areas of the quarry floor. Because the surface is parallel to bedding, widespread, and easily recognized in outcrop, it makes an excellent structural horizon for detailed mapping. One of the most common structures observed in the quarry floor are small domes and folds that are superimposed on the west-southwest regional dip. Typically, the domes are round to elliptical, 30 to 200 feet in diameter, and up to 20 feet high. Some of the domes are sites where the Mifflin limestone has undergone significant dissolution forming clay-filled cavities. The folds consist mainly of northwest-trending monoclines and anticlines that appear to have as much as 25 feet of relief and are hundreds of feet long.

Similar but smaller folds are exposed in the banks of Pine Creek and its tributaries in Sec. 15 and 16, T23N, R9E. In the SW<sup>1</sup>/<sub>4</sub> of Sec. 16, flat-lying rocks of the Quimbys Mill Formation are widely exposed in the floor of a quarry. At the north end of the quarry, the Quimbys Mill dips abruptly 5 degrees to the north-northeast beneath a small eastward-flowing tributary to Pine Creek. On the north side of the tributary is a small quarry exposure of the Dunleith Formation. The dip brings the Dunleith down to the same elevation as the Quimbys Mill.

A monocline is prominently exposed nearby in the banks of Pine Creek and its tributaries (NW SE Sec. 16). Here Galena strata have a northwest strike and dip about 25 degrees to the northeast. The fold trends in a southeasterly direction into the SW1/4 SW1/4 of Sec. 15. A second northwest-trending monocline is exposed in the west bank of Pine Creek in NW SE Sec. 15. Platteville strata dip about 20 degrees to the northeast. This monocline is on trend with and possibly continuous with a monocline exposed in a ravine across Pine Creek Valley about 2,000 feet to the west (NE NW SW Sec. 15). A concealed fault is inferred to be present north of and adjacent to this the monocline (center of Sec. 15).

A northwest-trending fault is well exposed in a prominent ravine on the north side of Rock River approximately ½ mile west of Grand Detour (SW NE Sec. 11, T22N, R9E). Here the St. Peter Sandstone is upthrown at least 30 feet in juxtaposition with dolomite beds of the Platteville Group.

Bedrock fracture patterns indicate dominant northwest and subdominant southwest trends similar to other parts of northern Illinois.

## **ECONOMIC RESOURCES**

High-calcium, low-magnesium limestone has been mined for cement production at St. Marys Cement Company in Dixon, Illinois (S<sup>1</sup>/<sub>2</sub> Sec. 16, SW<sup>1</sup>/<sub>4</sub> 15, S<sup>1</sup>/<sub>2</sub> NE<sup>1</sup>/<sub>4</sub> Sec. 21, and S<sup>1</sup>/<sub>2</sub> Sec. 22, T22N, R9E). The Mifflin Formation is the primary target, but high-quality limestone locally is present in the lower 10 to 20 feet of the overlying Grand Detour Formation as well. The limestone facies extends west of the St. Marys quarries through the bluffs of the Rock River and northward into the Pine Creek area. Approximately 10 feet of limestone is present in the Mifflin Formation in a core drilled near the intersection of Lowell Park Road and Penn Corner Road (SE NE NE Sec. 31, T23N, R9E; IP number 121412600200; ISGS core C15474). About 15 feet of Mifflin limestone is exposed in the banks of a tributary to Pine Creek (NE NW Sec. 34, T23N, R9E) and in another tributary 1,000 feet to the northeast (SW SE Sec. 27, T23N, R9E). North of here through the Grand Detour Quadrangle the limestone grades to dolomite. No limestone is present in the extensive Platteville outcrops in the northwest-trending ravine between Harmony and Columbian Roads (NW Sec. 14, T23N, R9E) or in the Burlington Northern Railroad cut (NW NW SE and NE SW Sec. 7, T23N, R10E).

## Dolomite

Limestone

In northern Illinois the Platteville and Galena dolomite is widely quarried for use as aggregate, road surfacing material, agricultural lime, and rip-rap. Active quarries in the Grand Detour Quadrangle include those on the north side of Dixon, Illinois (S<sup>1</sup>/<sub>2</sub> Sec. 16, SW<sup>1</sup>/<sub>4</sub> 15, S<sup>1</sup>/<sub>2</sub> NE<sup>1</sup>/<sub>4</sub> Section 21, and S<sup>1</sup>/<sub>2</sub> Sec. 22, T22N, R9E) and near the intersection of Judson and Anterior Roads. Commercial grade flagstone has also been mined from the latter.

#### Sandstone

The St. Peter Sandstone is actively being mined by the UNIMIN Corporation in the northeastern part of

glass, filter and molding sand, and as an abrasive. The St. Peter Sandstone is also used in hydrofracturing of oil and gas wells, as well as water wells, in order to increase flow rates. The St. Peter occurs at the bedrock surface in a wide area along the east side of the quadrangle.

# ACKNOWLEDGMENTS

We thank the numerous landowners who allowed us onto their land for outcrop study. Special thanks go to St. Marys Cement Company, which gave us access to their active and abandoned quarries, and to William G. Stuff for allowing us to drill a cored test hole on his family's property. Jacquelyn L. Hannah provided field assistance. Mary J. Seid provided valuable insight to the geology at the Burlington Northern Railroad cut and to the pre-Glenwood deformation observed in the nearby sandstone mines. Special thanks go to Donald E. Luman for the LiDAR maps and to Jane E. Johnshoy Domier for the graphic and cartographic layout and design.

This research was supported in part by the U.S. Geological Survey (USGS) National Cooperative Geologic Mapping Program (STATEMAP) under USGS award number G10AC00418. The views and conclusions contained in this document are those of the authors and should not be interpreted as necessarily representing the official policies, either expressed or implied, of the U.S. Government.

# REFERENCES

- Knappen, R.S., 1926, Geology and mineral resources of the Dixon Quadrangle: Illinois State Geological Survey, Bulletin 49, 141 p.
- Kolata, D.R., and T.C. Buschbach, 1976, Plum River Fault Zone of northwestern Illinois: Illinois State Geological Survey, Circular 491, 20 p.
- Kolata, D.R., T.C. Buschbach, and J.D. Treworgy, 1978, The Sandwich Fault Zone of northern Illinois: Illinois State Geological Survey, Circular 505, 26 p.
- Kolata, D.R., W.D. Huff, and S.M. Bergström, 1996, Ordovician K-bentonites of eastern North America: Geological Society of America, Special Paper 313, 84 p.
- Kolata, D.R., W.D. Huff, and S.M. Bergström, 1998, Nature and regional significance of unconformities associated with the Middle Ordovician Hagan K-bentonite complex in the North American midcontinent: Geological Society of America Bulletin, v. 110, no. 6, p. 723–739.
- Kolata, D.R., W.D. Huff, and S.M. Bergström, 2001, The Ordovician Sebree Trough: An oceanic passage to the Midcontinent United States: Geological Society of America Bulletin, v. 113, no. 8, p. 1067-1078.
- McGarry, C.S., 1999, Bedrock geology of Lee County, Illinois: Illinois State Geological Survey, Open File Series, OFS 1999-1e Lee BG, 1:62,500.
- Piskin, K., and S.M. Bergström, 1975, Glacial drift in Illinois: Thickness and character: Illinois State Geological Survey, Circular 490, 35 p.
- Seid, M.J., and D.R. Kolata, 2011, Newly discovered evidence for Late Ordovician structural deformation in Ogle County, Illinois: GSA Abstracts with Programs, Minneapolis, Paper no. 202-10, v. 45, no. 5, p. 493.
- Templeton, J.S., and H.B. Willman, 1952, Guidebook for the Sixteenth Annual Field Conference of the Tri-State Geological Society-Central Northern Illinois: Illinois State Geological Survey, 47 p.
- Templeton, J.S., and H.B. Willman, 1963, Champlainian Series (Middle Ordovician) in Illinois: Illinois State Geological Survey, Bulletin 89, 260 p.
- Weller, S., 1906, The geologic map of Illinois: Illinois State Geological Survey, Bulletin 1, 24 p.
- Willman, H.B., and D.R. Kolata, 1978, The Platteville and Galena Groups in northern Illinois: Illinois State Geological Survey, Circular 502, 75 p.
- Worthen, A.H., 1875, Geological map of Illinois: Geological Survey of Illinois, Geology and palaeontology, v. 6, 577 p.

the Grand Detour Quadrangle (Sec. 18, T23N, R10E). This sandstone is used for the manufacture of



Figure 2 North-facing highwall in a sandstone mine near the center of Sec. 18 (T23N, R10E) revealing an episode of Late Ordovician structural deformation. The St. Peter Sandstone is folded into a broad north-northeast-trending syncline and anticline. Structurally tilted up to 20 degrees, the St. Peter strata were subsequently eroded and covered by flat-lying shale, dolomite, and sandstone of the Glenwood Formation, forming a pronounced angular unconformity. The center of the highwall is approximately 100 feet high.



# IGQ Grand Detour-BG Sheet 2 of 2