


SURFICIAL GEOLOGY OF MAPLE PARK QUADRANGLE  
KANE AND DE KALB COUNTIES, ILLINOIS

Department of Natural Resources  
ILLINOIS STATE GEOLOGICAL SURVEY  
William W. Shills, Chief

Illinois Preliminary Geologic Map  
IPGM 5 Maple Park-SG

David A. Grimley  
2004



QUATERNARY DEPOSITS

HUDSON EPISODE (~12,000 years before present (B.P.) to today)

gp Grayslake Peat

Fibrous peat, muck, organic silt and clay; interbedded with sand, silt, and clay in some places, up to 20 feet thick; intertongues with Equality Formation and Cahokia Formation; may overlie Cahokia, Equality or Henry Formations.

c Cahokia Formation

Stratified to massive sand, silt, and clay; generally less than 10 feet thick; dominantly silty redeposited loess and lacustrine sediment; occurs in floodplains and channels of modern streams and streams converted to drainage ditches. Alluvium.

WISCONSIN EPISODE (~55,000 - 12,000 years B.P.)  
MASON GROUP (sorted sediments)

e Equality Formation

Laminated to massive clay and silt, containing some fine to medium sand beds; 5 to 25 feet thick; intertongues with other units in Mason and Wedron Groups; occurs in proglacial, supraglacial, slackwater, and some modern lake basins; commonly underlain by Tiskilwa Formation, Batesown Member, Lemont Formation, or Henry Formation. Lacustrine sediment.

h Henry Formation (except Wasco facies)

Stratified to massive sand and gravel containing beds of silt, clay, and diamictic; generally well-sorted, cross-bedded to plane-bedded; up to 60 feet thick; intertongues with Equality Formation and Wedron Group units; intertongues or occurs under Batesown Member, Lemont Formation, occurs extensively under the Tiskilwa Formation; deposited in glacial meltwater channels, outwash plains, deltas, bars. Outwash.

hw Wasco facies, Henry Formation

Irregularly bedded and moderately sorted sand and gravel, containing lenses of silt, clay, and diamictic; 5 to 30 feet thick; associated with Batesown Member of Wedron Group; may contain a covering of 2 to 10 feet of loose loamy ablation till; occurs in kames in the Arlington Moraine in the southern third of the map. Ice-contact and ice-marginal sediment.

WISCONSIN EPISODE (~55,000 - 12,000 years B.P.)  
WEDRON GROUP (diamictic units)

lb Batesown Member, Lemont Formation

Silt loam to loam diamictic, gray to gray-brown, oxidizing to yellow-brown; as much as 35 feet thick; upper portion may be mixed with stratified and interbedded silt and sand; lower portion is more likely to be massive subglacial till. Till and ice-marginal sediment.

t Tiskilwa Formation

Loam to clay loam diamictic; pink to reddish-brown to gray; locally contains thick beds of sand and gravel; sometimes stratified in upper portions; occurs at the surface in the northwest and northeast portions of the quadrangle, where it is up to 150 feet thick. Till and ice-marginal sediment.

ILLINOIS EPISODE (~200,000 - 130,000 years B.P.)


g Glasford Formation (in cross sections only)

Loam to clay loam diamictic, pinkish-brown to brown to gray; up to 110 feet thick; contains some beds of sorted sediment, within and especially at the base of the unit. Till and ice-marginal sediment and channel deposits.

Map Legend (data points and lines)

- Outcrops, exposures or hand augers (described by ISGS)
- Stratigraphic test hole (drilled by ISGS)
- Engineering boring
- Water well boring
- Line of cross section

note: data point labels for borings indicate the 5-digit "county number" which is a portion of the 12-digit API number available at the Geologic Records Unit of the Illinois State Geological Survey; outcrops and hand augers have field numbers indicated



INTRODUCTION

This map of surficial (Quaternary) deposits for the Maple Park Quadrangle is located in northeastern Illinois about 50 miles west of Chicago. Studied deposits include unconsolidated sediments down to bedrock. The map is intended to provide an important framework for land and groundwater use, engineering assessment, economic development and geological studies in the area. This study is part of a broader geologic mapping program undertaken by the ISGS for 7.5-minute quadrangles in the Chicago Metro Region and was partially funded by a USGS-STATEMAP contract.

Glacial sediment, from 70 to 270 feet thick in this quadrangle, was deposited during successive advances of glaciers during both the Illinois and Wisconsin Episodes. At least one advance of Illinois Episode glaciers and two advances of Wisconsin Episode glaciers have deposited sediments in this part of western Kane and eastern De Kalb Counties (Curry et al., 1999; Grimley and Curry, 2001). This quadrangle is unusual for Illinois in that Lake Michigan Lobe ice advanced from a southeasterly direction into this area during the last glaciation (Fig. 1). The underlying bedrock, Paleozoic carbonates and shales, are not known to outcrop in the Maple Park 7.5' Quadrangle.

SURFICIAL DEPOSITS

Illinois Episode till and sand and gravel (both Glasford Formation in cross sections) are preserved primarily in bedrock lowlands or buried valleys and at elevations below about 750 to 800 feet. No outcrops of Illinois Episode deposits are known in the Maple Park quadrangle; however, they were encountered in several stratigraphic test borings and many water wells. The upper 5 to 10 feet of Glasford Formation deposits were intensely weathered during the Sangamon interglacial episode (from about 135,000 to 55,000 years before present). This physically and chemically weathered zone, known as the Sangamon Gossol, is often described as a "green clay" in water well drillers' logs. The Sangamon Gossol is an excellent marker bed for separation of Illinois Episode and Wisconsin Episode deposits (Curry, 1989).

During the Wisconsin Episode, a pink loam to clay loam diamictic with some sand and gravel bodies, classified as Tiskilwa Formation, was deposited in all areas of the quadrangle. The Tiskilwa Formation is up to 150 feet thick and is generally a dense and uniform subglacial till. This unit is distinct from other Wedron Group units in its color, texture, clay mineralogy, and engineering properties (Wickham et al., 1988; Curry et al., 1999). The Tiskilwa Formation was first deposited in the Marengo Moraine, whose western flank is in the northeastern portion of the Maple Park Quadrangle (Fig. 1), between about 25,000 and 22,000 radiocarbon years ago (Hansel and Johnson, 1996; Curry et al., 1999). Following this, a second phase of Tiskilwa Formation deposition occurred over much of the quadrangle as a result of ice advances (from south to north; Fig. 1) which formed the Bloomington Moraine System between about 22,000 and 19,000 radiocarbon years ago (Hansel and Johnson, 1996; Curry et al., 1999). The Tiskilwa Formation is thickest in the northeastern portion of this quadrangle probably since both phases of deposition occurred here. In the north-central portion of this quadrangle, the Tiskilwa Formation is overlain by thin lacustrine deposits (Equality Formation) and, in places, underlain by thin sand and gravel outwash (Ashmore Tongue, Henry Formation) that may represent proglacial deposits of the Marengo Moraine ice front (A-A' and B-B' cross-sections).

In the southern one-third of the quadrangle, a gray to yellow-brown, silt loam to loam diamictic (Batesown Member, Lemont Formation) is found in the Arlington Moraine (Fig. 1) and overlying the Tiskilwa Formation. The Batesown diamictic is either, less pink, and contains slightly more silt than the Tiskilwa diamictic. The Arlington Moraine marks a readvance position of the ice as it advanced from south to north during overall retreat of the Lake Michigan Lobe. Sand and gravel (Henry Formation) is sometimes found between the Batesown and Tiskilwa diamictics or in kame hills (Wasco facies, Henry Formation) on the moraine (C-C' cross-section).

A large area of lacustrine sediment exists north of the Arlington Moraine as ice advancing from the south caused meltwater to pond between the glacier and the Bloomington Moraine System. Stratified fine sand, silt, and clay (Equality Formation) occurs as a veneer, up to 25 feet thick, in north-central portions of the lake (C-C' and B-B' cross-sections). Other areas of lake sediment occur on both the Arlington or Bloomington Moraines. Many coarser sand bars and deltas in the glacial lake, as well as outwash below lake sediment, are mapped as Henry Formation. Sand and gravel outwash up to 60 feet thick occurs in front of the Arlington Moraine. The upper portion of this sediment is likely proglacial to this moraine, but some may be related to older advances.

The Grayslake Peat, up to 20 feet thick, is common in depressions and low-lying areas, within and adjacent to lake and outwash plains in this landscape. During postglacial times, peat and organic silts were deposited in swampy depressions and preserved due to anoxic conditions. Modern (postglacial) stream sediment, primarily sand and silt deposits, is inset into older glacial deposits along the lowlands adjacent to Union and Virgil Ditches. Being far from the base level influence of major rivers and having had only about 18,000 years since glaciation for landscape development (a short time geologically speaking), these stream channels have not yet developed well defined valleys or caused significant dissection. Constructional landforms (moraines, kames, lake plains, kettle holes) dominate the landscape in this quadrangle.

CROSS SECTIONS

Sand and gravel bodies are stippled on cross sections only where data indicates their presence. Additional sand and gravel lenses undoubtedly occur within the Glasford, Tiskilwa, and Batesown tills. A two- to four-foot thick cover of loess at the ground surface is not shown in the cross sections, nor are most other geologic units that are less than 5 feet in maximum thickness. Water wells and test holes used for the three cross sections are mainly transported from within 2000 feet of the cross section lines (see map). Those data points transported from ~2000 feet are indicated as dashed vertical lines in cross sections (cross-section A-A' only). Data points were transported to positions on the cross section with similar geomorphology and surface elevations. Many water wells extend deep into bedrock and so their full extent is not always shown.

DATASOURCES

This surficial geologic map is based in part upon soil series parent materials compiled from the Soil Survey of Kane County (Goldard, 1979; scale 1:15,840) and De Kalb County (Hinkley, 1978; scale 1:15,840), but was considerably modified based upon field observations and new drill cores obtained as part of this STATEMAP project.




Figure 1. Wisconsin Episode moraines in the Lake Michigan Lobe of northeastern Illinois; modified from Wilman and Frye (1970). Kane County is outlined in black.

Cross Section Legend

- Sand and gravel, may contain silt beds
- Diamictic, massive silt, or other fine-grained sediment
- Subsurface boring used for
- Subsurface boring used for

Surface contaminants provide a potential threat to groundwater supplies in near-surface aquifers that are not overlain by a confining (clayey, unfactured) aquitard, such as clayey till deposits or lake deposits (Berg, 1984). Shallow sand and gravel aquifers, such as Henry Formation exposed at the surface or buried by a thin loess cap (< 4 feet) are most vulnerable to agricultural or industrial contaminants. Tiskilwa diamictic (contains an average of 25-30 % clay) is an excellent aquifer when it is uniform and does not contain sand bodies within it. Batesown diamictic (typically 15-20 % clay) is a less reliable aquifer because this till is less clayey than Tiskilwa diamictic and it also is generally thinner and more heterogeneous, containing numerous sand bodies and lenses.

REFERENCES

- Anderson, R.C., 1964. Sand and gravel resources of De Kalb County. Illinois State Geological Survey, Circular 367, 16 p.
- Berg, R.C., Kempton, J.P., Cartwright, K., 1984. Potential for contamination of shallow aquifers in Illinois. Illinois State Geological Survey, Circular 532, 30 p.
- Block, D.A., 1960. Sand and gravel resources of Kane County. Illinois State Geological Survey, Circular 299, 11 p.
- Curry, B.B., Grimley, D.A., and J.A. Stivers, 1999. Quaternary geology, geomorphology, and climatic history of Kane County, Illinois. Illinois State Geological Survey, Geobasebook 28, 44 p.
- Curry, B.B., Giese, A.M., Hansel, M.J., Vaiden, R.C., Bauer, R.A., Schumacher, D.A., Norton, K.A., Dixon, J.E., W.G., 1988. Geological-geomorphological index for using the Superconducting Super Collider in Illinois: Results of the 1986 ice drilling program. Illinois State Geological Survey, Environmental Geology Notes 122, 188 p.
- Curry, B.B., 1989. Absence of Algonquin glaciation in Illinois. Quaternary Research, v. 31, p. 1-13.
- Goldard, T.M., 1979. Soil survey of Kane County, Illinois. USDA Soil Conservation Service, 179 p.
- Goldman, H.B., 1984. Sand and gravel, in Donald D. Carr, ed., Industrial minerals and rocks, 6th Edition: Lithology, Colorado. Society for Mining, Metallurgy, and Exploration, Inc., p. 969-977.
- Grimley, D.A., and Curry, B.B., 2001. Surficial Geologic Map, Elmhurst Quadrangle, Kane County, IL. Illinois State Geological Survey, Illinois Geologic Quadrangle Map (IGQ) Elmhurst-SG, scale 1:24,000. Chicago, D.L., 1969. Glacial geology of Kane County, Illinois. University of Illinois at Urbana-Champaign, Ph.D. dissertation, 211 p.
- Hansel, A.K., and Johnson, W.H., 1996. Wedron and Mason Groups: Lithostratigraphic reclassification of deposits of the Wisconsin Episode, Lake Michigan Lobe. Illinois State Geological Survey, Bulletin 104, 116 p.
- Hinkley, K.C., 1978. Soil survey of De Kalb County, Illinois. University of Illinois at Urbana-Champaign Agricultural Experiment Station, 69 p.
- Leverett, F., 1899. The Illinois Glacial Lake. U.S. Geological Survey, Monograph 38, 817 p.
- Wickham, S.S., W. H. Johnson, and H.D. Glas. 1988. Regional geology of the Tiskilwa Till Member, Wedron Formation, northeastern Illinois. Illinois State Geological Survey, Circular 543, 35 p.
- Wilman, H.B. and J.C. Frye, 1970. Pleistocene stratigraphy of Illinois. Illinois State Geological Survey, Bulletin 94, 204 p.

ACKNOWLEDGMENTS

Discussions with B.B. Curry and A.K. Hansel were valuable for the development of this map. T.J. McElroy and D. Collins helped considerably with computerization of the map in ARC/INFO and Arcview. J. Danner helped to design the layout and base map materials. H.D. Glas supplied important clay mineralogical data, used to aid correlation of till units.